

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

1012E-000300US

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/868968
Not yet knownINTERNATIONAL APPLICATION NO.
PCT/GB99/04382INTERNATIONAL FILING DATE
23 December 1999PRIORITY DATE CLAIMED
23 December 1998TITLE OF INVENTION TREATMENT AND DIAGNOSIS OF CANCER USING INOSITOLPHOSPHOGLYCANS
ANTAGONISTS

APPLICANT(S) FOR DO/EO/US

RADEMACHER, Thomas William

532 Rec'd PCT/PTO 22 JUN 2001

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
 2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
 3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
 4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
 5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
 6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
 7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
 8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
 9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
 10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).
- Items 11. to 16. below concern document(s) or information included:
11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
 12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
 13. ☒ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
 14. ☐ A substitute specification.
 15. ☐ A change of power of attorney and/or address letter.
 16. ☒ Other items or information:
 - Verified Statement of Small Entity Status
 - Copy of IPER with Amendment under Article 34
 - Receipt Acknowledgement postcard
 - Certificate of Mailing

U.S. APPLICATION NO. (If known, see 37 CFR 1.51)

09/868968

INTERNATIONAL APPLICATION NO.

PCT/GB99/04382

ATTORNEY'S DOCKET NUMBER

1012E-000300US

17. ☒ The following fees are submitted:**BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) :**

Neither international preliminary examination fee (37 CFR 1.482)
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO
and International Search Report not prepared by the EPO or JPO \$970.00

International preliminary examination fee (37 CFR 1.482) not paid to
USPTO but International Search Report prepared by the EPO or JPO. \$840.00

International preliminary examination fee (37 CFR 1.482) not paid to USPTO but
international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$760.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)
but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$670.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)
and all claims satisfied provisions of PCT Article 33(1)-(4) \$96.00

ENTER APPROPRIATE BASIC FEE AMOUNT =**CALCULATIONS** PTO USE ONLY

\$ 840.00

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(e)).

\$

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	22 - 20 =	2	X \$18.00
Independent claims	4 - 3 =	1	X \$80.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$270.00
TOTAL OF ABOVE CALCULATIONS =			\$ 956.00
Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).			\$ 478.00
SUBTOTAL =			\$ 478.00
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).			\$
TOTAL NATIONAL FEE =			\$ 478.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property			\$
TOTAL FEES ENCLOSED =			\$ 478.00
			Amount to be:
			refunded
			charged
			\$ 478.00

\$ 36.00

\$ 80.00

\$

\$ 956.00

\$ 478.00

\$ 478.00

\$

\$ 478.00

\$

\$ 478.00

Amount to be:

refunded

charged

\$

\$ 478.00

a. ☐ A check in the amount of \$_____ to cover the above fees is enclosed.b. ☒ Please charge my Deposit Account No. 50-0893 in the amount of \$ 478.00 to cover the above fees. A duplicate copy of this sheet is enclosed.c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 50-0893. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Emily M. Haliday
LAW OFFICES OF JONATHAN ALAN QUINE
P.O. BOX 458
Alameda, CA 94501
United States of America

SIGNATURE

Emily M. Haliday

NAME

38,903

REGISTRATION NUMBER

Atty. Docket No. 1012E-000300US

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
(37 CFR 1.9(d) & 1.27(c)) - SMALL BUSINESS CONCERN

Applicant or Inventor: Thomas William Rademacher and Hugo CaroApplication or Patent No.: U.S. National Phase Application of PCT/GB99/04382Filed or Issued: December 23, 1999 (International Filing Date)Title: TREATMENT AND DIAGNOSIS OF CANCER USING INOSITOLPHOSPHOGLYCANS ANTAGONISTS

I hereby declare that I am:



the owner of the small business concern identified below

an official of the small business concern empowered to act on behalf of the concern identified below.

Name of Small Business Concern: Rademacher Group LimitedAddress of Small Business Concern: 6 St. Andrew Street, London EC4A 3JX, United Kingdom

I hereby declare that the above-identified small business concern qualifies as a small business concern as defined in 13 CFR 121.12, and reproduced in 37 CFR 1.9(d), for purposes of paying reduced fees to the United States Patent and Trademark Office, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention, entitled **TREATMENT AND DIAGNOSIS OF CANCER USING INOSITOLPHOSPHOGLYCANS ANTAGONISTS** by inventor(s) Thomas William Rademacher and Hugo Caro described in:



the specification filed herewith,

Attorney Docket No. 1012E-000300US, filed 6/22/2001.

Patent No. _____, issued _____.

If the rights held by the above identified small business concern are not exclusive, each individual, concern or organization having rights in the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person made the invention, or by any concern that would not qualify as a small business concern under 37 CFR 1.9(d), or a nonprofit organization under 37 CFR 1.9(e).

*NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

Name _____

Address _____



Individual



Small Business Concern



Nonprofit Organization

Name _____

Address _____



Individual



Small Business Concern



Nonprofit Organization

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b)).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and fee like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

Name of Person Signing: THOMAS RADEMACHERTitle of Person if Other than Owner: DIRECTORAddress of Person Signing: Rademacher Group Limited, 6 St. Andrew Street, London EC4A 3JX, United KingdomSignature Thomas RademacherDate 20/26/01

I hereby certify that this correspondence is being deposited with the United States Postal Service express mail in an envelope addressed to:
Assistant Commissioner for Patents Box PCT, Washington, D.C. 20231, on

6/22/01

LAW OFFICES OF JONATHAN ALAN QUINE

By:

Debrah Berwick

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Thomas William Rademacher; Hugo Caro

Examiner: Unknown

Application No.: Unknown

Art Unit:

Filed: 6/22/2001

For: **Treatment and Diagnosis of Cancer
Using Inositolphosphoglycans Antagonists****PRELIMINARY AMENDMENT**

Box PCT
Assistant Commissioner for Patents
Washington, D.C. 20231

Dear Sir:

Prior to calculating the claim fees, please amend the claims of the above-referenced application as indicated below.

In the Claims:

Please amend the claims by canceling claims 1-21 without prejudice and adding the following claims:

22. A method of reducing tumor cell proliferation comprising contacting a tumor cell with a substance which is an inositolphosphoglycan (IPG) antagonist having the property of reducing tumor cell proliferation.

23. The method of claim 22, wherein the tumor cell is in vivo in a subject and the substance is administered to the subject for the treatment of cancer.

24. The method of claim 22, wherein the IPG antagonist is:

(a) a substance which is capable of inhibiting the release of IPGs; or,

(b) a substance capable of reducing the levels of IPGs by binding to the IPGs; or,

(c) a substance which is a competitive agent which is capable of reducing an effect of IPGs.

25. The method of claim 24, wherein the IPG antagonist is a competitive IPG antagonist.

26. The method of claim 24, wherein the IPG antagonist is an anti-IPG antibody which is capable of specifically binding IPGs.

27. The method of claim 26, wherein the antibody is capable of neutralizing an activity of the IPGs.

28. The method of claim 27, wherein activity of the IPGs is the proliferation of tumor cells.

29. The method of claim 26, wherein the antibody is a monoclonal antibody produced by hybridoma 2F7, 2D1 or 5H6, deposited at ECACC under accession numbers 98051201, 98031212 and 98030901, respectively.

30. The method of claim 24, wherein the antagonist is an inhibitor of glycosylphosphatidylinositol specific phospholipase type C (GPI-PLC).

31. A method for the diagnosis and/or prognosis of cancer, the method comprising determining the presence or amount of inositolphosphoglycans in a sample from a patient.

32. The method of claim 31, wherein the presence or amount of the IPGs is causing tumor cell proliferation.

33. The method of claim 31, wherein the method comprises:

(a) contacting a sample from a patient with a solid support having immobilized thereon a binding agent having a binding site which is capable of specifically binding to the IPGs under conditions in which the IPGs bind to the binding agent; and,

(b) determining the presence or amount of the IPGs bound to the binding agent.

34. The method of claim 33, wherein step (b) comprises

(i) contacting the solid support with a developing agent which is capable of binding to occupied binding sites, unoccupied binding sites or the bound IPGs, the developing agent comprising a label, and (ii) detecting the label to obtain a value representative of the presence or amount of the IPGs in the sample.

35. The method of claim 34, further comprising comparing the value with standards from healthy or cancerous tissues.

36. The method of claim 34, wherein the label is selected from the group consisting of a radioactive label; a chemiluminescent label; a fluorophor; a phosphor; a laser dye; a chromogenic dye; a macromolecular colloidal particle; a latex bead which is coloured, magnetic or paramagnetic; and an enzyme which catalyses a reaction producing a detectable result.

37. The method of claim 33, wherein the binding agent immobilized on the solid support is an antibody which is capable of binding to the IPGs.

38. The method of claim 33, wherein the binding agent is immobilized at a predefined location on the solid support.

39. A method for purifying or isolating a P or A-type substance, wherein the substance is a cyclitol containing carbohydrate which is:

(i) a P-type substance having the biological activity of activating pyruvate dehydrogenase (PDH) phosphatase; or,

(ii) an A-type substance having the biological activity of inhibiting cAMP dependent protein kinase,

wherein the method comprises contacting a sample containing P or A-type substance with microcrystalline cellulose.

40. The method of claim 39, wherein the method involves contacting a sample containing P or A-type substance with a column containing cellulose and eluting the substance from the column.

41. A method of purifying or isolating a P or A-type substance, wherein the substance is a cyclitol containing carbohydrate which is:

(i) a P-type substance having the biological activity of activating pyruvate dehydrogenase (PDH) phosphatase; or,

(ii) an A-type substance having the biological activity of inhibiting cAMP dependent protein kinase; wherein the method comprises:

(a) loading a column containing microcrystalline cellulose with a sample containing the P or A-type substance so that P or A-type substance binds to the column; and,

(b) eluting the P or A-type substance from the column.

42. The method of claim 41, further comprising the step of dissolving the sample containing the P or A-type substance in 4/1/1 butanol/water/ethanol (B:W:E) prior loading on the column.

43. The method of claim 41, further comprising the step of washing the column with B:W:E and methanol.

These amendments are made without prejudice and are not to be construed as abandonment of the previously claimed subject matter.

REMARKS

Claims 22-43 are pending with entry of this amendment, claims 1-21 being cancelled and claims 22-43 being added herein. These amendments introduce no new matter.

Prompt examination of the above claims is respectfully requested. If a telephone conference would expedite prosecution of this application, the Examiner is invited to telephone the undersigned at (510) 337-8894.

LAW OFFICES OF JONATHAN ALAN QUINE
P.O. BOX 458
Alameda, CA 94501
Tel: 510 337-7871
Fax: 510 337-7877

Respectfully submitted,



Emily M. Haliday
Reg. No: 38,903

TREATMENT AND DIAGNOSIS OF CANCER USING INOSITOLPHOSPHOGLYCANS ANTAGONISTSField of the Invention

5 The present invention relates to materials and methods for the treatment and diagnosis of cancer.

Background of the Invention

10 It has been observed that when tumour cells are grown in culture, the rate of proliferation of the cells depends on the cell density in the culture. It has been hypothesised that this increase in the rate of proliferation as the number of cells increases is due to the presence of tumour autocrine factors (TAFs) in the cell supernatant. TAFs are believed to be growth factors produced by the tumour cells which cause them to proliferate in the absence of external stimuli such as hormones.

15
20 There is some experimental evidence supporting the existence of TAFs in the culture supernatant. If the supernatant from a high density tumour cell culture is applied to a dilute culture, the rate of proliferation of the cells in the dilute culture quickly accelerates to the level associated with cells in a high density culture. It has also been observed that if the supernatant is boiled prior to addition to the dilute culture, the acceleration in the proliferation rate of the cells is not observed.

25
30 Witter et al (1987) have reported that conditioned medium from Reuber H-35 or Fao hepatoma cells contained autocrine factors that both stimulated DNA synthesis and activated acetyl-coenzyme A (CoA) carboxylase in serum deprived cells. They also found that the factors increased the cell number and mitotic index in tumour

35

cell cultures. The investigators fractionated conditioned medium from H-35 hepatoma cells and found that two low molecular weight components (MW < 1000) co-purified with the activity of increasing DNA synthesis. Radiolabelling incorporation studies suggested the possible incorporation of glucosamine and galactose, but no incorporation of *myo*-inositol or mannose. However, the authors were unable to purify, isolate or characterise the autocrine factor from conditioned medium and do not provide any definite suggestions as to the structure or activity of their autocrine factor.

Summary of the Invention

Broadly, the present invention is based on the isolation and identification of TAFs, and in particular on the finding that TAFs are inositolphosphoglycans (IPGs) such as A or P-type IPGs. In particular, the results show that the TAFs include *myo*-inositol and by extension A-type IPGs.

Accordingly, in a first aspect, the present invention provides the use of a substance which is an IPG antagonist for the preparation of a medicament for the treatment of cancer. Preferably, the IPG antagonist has the property of reducing tumour cell proliferation.

In the present application, "IPG antagonists" includes substances which have one or more of the following properties:

(a) substances which are capable of inhibiting release of IPGs, e.g. inhibitors of enzymes which cause their release such as glycosylphosphatidylinositol specific phospholipases (GPI-PL);

(b) substances which are capable of reducing the levels of IPGs by binding to them, e.g. anti-IPG

antibodies, and preferably neutralising one or more IPG biological activities, more especially the activity of IPGs causing tumour cell proliferation; and/or,

(c) substances which are capable of reducing the effects of IPGs, e.g. competitive antagonists such as IPGs which are biologically inactive in the species in which they are used, e.g. the use of porcine liver A-type IPGs in humans.

In preferred embodiments, the IPG antagonist is an antibody capable of specifically binding IPGs and preferably which does not cross-react with the common reactive determinant of GPI anchored proteins. Preferably, the antibody is a monoclonal antibody, examples of which are disclosed below.

In other embodiments, the antagonist inhibits the release of IPGs. This suggests that an enzyme responsible for the release of the IPG TAFs is a GPI-PLC, which has not been previously identified in mammalian tissues. Therefore, inhibitors of GPI-PLC may be used as IPG antagonists for treatment of cancer.

In a further aspect, the present invention provides a method for the diagnosis or prognosis of cancer, the method comprising determining the presence or amount of IPGs in a sample from a patient. The presence or amount of IPGs in a given sample is typically then compared to standards from healthy and cancerous tissues.

Preferably, the presence or amount of the IPGs is determined by measuring a characteristic activity of an A or P-type IPG, details of which are set out below. Alternatively, the presence or amount of a characteristic component of IPGs and in particular IPG TAFs, could be

used as a diagnostic marker. Other components of IPGs that might be used as diagnostic markers are set out below.

5 In a further aspect, the present invention provides the use of cellulose column chromatography in the purification of IPGs.

10 In a further aspect, the present invention provides a method of purifying or isolating IPGs, the method comprising making contacting a sample containing IPGs with a column containing cellulose, and eluting the IPGs from the column. Preferred conditions preparing the column and eluting IPG containing fractions are set out below.

15 Embodiments of the present invention will now be described by way of example and not by limitation with reference to the accompanying figures.

Brief Description of the Figures

20 Figure 1 shows the fractionation of conditioned medium by cellulose column chromatography. The results are presented as counts per minute (cpm) per fraction. The arrows indicate the solvent system used at a particular fraction.

25 Figure 2 shows the effects on DNA synthesis of conditioned medium. Conditioned medium at two different concentrations (10% and 20%) was assayed for stimulating activity in test Fao cells. The results are presented as radioactivity (cpm) obtained after incorporation of ³H-thymidine into DNA, for control (no serum addition), FCS (5% foetal calf serum + 5% foetal bovine serum addition), and the fractions obtained. The arrows indicate the

30

35

eluent system used at a particular fraction.

Figure 3 shows phosphate analysis of conditioned medium fractionated by cellulose column chromatography. Results are presented as OD obtained per fraction. Arrows indicate the eluent system used at a particular fraction.

Figure 4 shows the effect on the proliferation of FaO cells of conditioned medium treated with anti-IPG monoclonal antibodies. Results are presented as radioactivity (cpm) obtained after incorporation of ^3H -thymidine into DNA, for control (no serum addition), FCS (5% foetal calf serum + 5% foetal bovine serum addition), and samples containing the antibodies.

Figures 5 and 6 show rat liver IPGs A and P fractionation profiles. The results are shown as radioactivity (cpm) obtained after incorporation of ^3H -thymidine into DNA of 3T3EFGTR17 cells, for control (no serum addition), FCS (10% foetal calf addition), and fractions. All fractions were assayed at a final IPG-A concentration of 1/80, except for fraction 26 of IPG-P (Fig. 5) which was also tested at a final concentration of 1/40 and 1/160. All fractions of the IPG-A (Fig. 6) were tested at a final concentration of 1/80. The arrows indicate the eluent system used at a particular fraction.

Figure 7 shows the phosphate analysis of rat liver IPG-P fractionated from rat liver by cellulose column chromatography. Results are presented as OD obtained per fraction. Arrows indicate the eluent system used at a particular fraction.

Detailed Description of the Invention

IPGs

Studies have shown that A-type mediators modulate the activity of a number of insulin-dependent enzymes such as cAMP dependent protein kinase (inhibits), adenylate cyclase (inhibits) and cAMP phospho-diesterases (stimulates). In contrast, P-type mediators modulate the activity of insulin-dependent enzymes such as pyruvate dehydrogenase phosphatase (stimulates), glycogen synthase phosphatase (stimulates) and cAMP dependent protein kinase (inhibits). The A-type mediators mimic the lipogenic activity of insulin on adipocytes, whereas the P-type mediators mimic the glycogenic activity of insulin on muscle. Both A-and P-type mediators are mitogenic when added to fibroblasts in serum free media. The ability of the mediators to stimulate fibroblast proliferation is enhanced if the cells are transfected with the EGF-receptor. A-type mediators can stimulate cell proliferation in the chick cochleovestibular ganglia.

Soluble IPG fractions having A-type and P-type activity have been obtained from a variety of animal tissues including rat tissues (liver, kidney, muscle brain, adipose, heart) and bovine liver. A- and P-type IPG biological activity has also been detected in human liver and placenta, RBC infected with *Plasmodium yoelii* and mycobacteria. The ability of an anti-inositolglycan antibody to inhibit insulin action on human placental cytotrophoblasts and BC3H1 myocytes or bovine-derived IPG action on rat diaphragm and chick cochleovestibular ganglia suggests cross-species conservation of many structural features.

A-type substances are cyclitol-containing carbohydrates, also containing Zn^{2+} ion and optionally phosphate and having the properties of regulating lipogenic activity

and inhibiting cAMP dependent protein kinase. They may also inhibit adenylate cyclase, be mitogenic when added to EGF-transfected fibroblasts in serum free medium, and stimulate lipogenesis in adipocytes.

P-type substances are cyclitol-containing carbohydrates, also containing Mn^{2+} and/or Zn^{2+} ions and optionally phosphate and having the properties of regulating glycogen metabolism and activating pyruvate dehydrogenase phosphatase. They may also stimulate the activity of glycogen synthase phosphatase, be mitogenic when added to fibroblasts in serum free medium, and stimulate pyruvate dehydrogenase phosphatase.

Methods for obtaining A-type and P-type IPGs are set out in Caro et al, 1997, and in WO98/11116 and WO98/11117. In summary, the methods disclosed in these applications involve:

(a) making an extract by heat and acid treatment of IPG source material;

(b) after centrifugation and charcoal treatment, allowing the resulting solution to interact overnight with an AG1-X8 (formate form) anion exchange resin;

(c) collecting a fraction having A-type IPG activity obtained by eluting the column with 50 mM HCl and/or collecting a fraction having P-type IPG activity obtained by eluting the column with 10mM HCl;

(d) adjusting to pH 4 (not to exceed pH 7.8) and lyophilising the fraction to isolate the substance;

(e) employing descending paper chromatography using 4/1/1 butanol/ethanol/water as solvent;

(f) purification using high-voltage paper electrophoresis in pyridine/acetic acid/water; and,

(g) purification using Dionex anion exchange chromatography or purification and isolation using Vydac

HPLC chromatography to obtain the isolated IPG.

As disclosed herein, it is also possible to employ column chromatography using cellulose, and especially microcrystalline cellulose, in the isolation or purification of IPGs. Exemplary conditions are provided in the experimental section below.

Antagonists

In the present invention, "IPG antagonists" includes substances which have one or more of the following properties:

(a) substances which are capable of inhibiting release of IPGs, e.g. inhibitors of enzymes which cause their release such as GPI-PL;

(b) substances which are capable of reducing the levels of IPGs by binding to them and preferably neutralising an IPG biological activity; and/or,

(c) substances which are capable of reducing the effects of IPGs, e.g. competitive antagonists such as IPGs which are biologically inactive in the species in which they are used, e.g. the use of A-type IPGs as obtainable from porcine liver in humans.

Under (b) and (c), preferably the biological activity of the IPGs which is affected by the antagonists is the tumour cell proliferation caused by the IPGs when they are added to cell cultures. This can be readily determined using the assays described in the examples or other techniques well known to those skilled in the art.

Examples of IPG antagonists of type (b) include specific binding proteins, e.g. naturally occurring specific binding proteins that can be obtained by screening biological materials for substances that bind to IPGs.

In a further example, the antagonists are antibodies capable of specifically binding to IPGs. The production of polyclonal and monoclonal antibodies is well established in the art. Examples of anti-IPG monoclonal antibodies are produced by hybridoma cell lines 2F7, 2D1 and 5H6 deposited at European Collection of Cell Cultures (ECACC) under accession numbers 98051201, 98031212 and 98030901 on 12 May 1998 ('201) and 9 March 1998 ('212 and '901).

Monoclonal antibodies are particularly useful as they can be subjected to the techniques of recombinant DNA technology to produce other antibodies or chimeric molecules which retain the specificity of the original antibody. Such techniques may involve introducing DNA encoding the immunoglobulin variable region, or the complementarity determining regions (CDRs), of an antibody to the constant regions, or constant regions plus framework regions, of a different immunoglobulin. See, for instance, EP 0 184 187 A, GB 2 188 638 A or EP 0 239 400 A. A hybridoma producing a monoclonal antibody may be subject to genetic mutation or other changes, which may or may not alter the binding specificity of antibodies produced.

Antibodies may be obtained using techniques which are standard in the art. Methods of producing antibodies include immunising a mammal (e.g. mouse, rat, rabbit, horse, goat, sheep or monkey) with an immunogen, e.g. an IPG or a fragment thereof. Antibodies may be obtained from immunised animals using any of a variety of techniques known in the art, and screened, preferably using binding of antibody to antigen of interest. For instance, Western blotting techniques or immunoprecipitation may be used (Armitage et al, Nature,

357:80-82, 1992). Isolation of antibodies and/or antibody-producing cells from an animal may be accompanied by a step of sacrificing the animal.

5 As an alternative or supplement to immunising a mammal with an IPG, an antibody specific for the IPG may be obtained from a recombinantly produced library of expressed immunoglobulin variable domains, e.g. using lambda bacteriophage or filamentous bacteriophage which
10 display functional immunoglobulin binding domains on their surfaces; for instance see WO92/01047.

Antibodies according to the present invention may be modified in a number of ways. Indeed the term "antibody"
15 should be construed as covering any binding substance having a binding domain with the required specificity. Thus, antagonist antibodies includes antibody fragments, derivatives, functional equivalents and homologues of antibodies, including synthetic molecules and molecules
20 whose shape mimics that of an antibody enabling it to bind an antigen or epitope.

Examples of antibody fragments which capable of binding an IPG or other binding partner are the Fab fragment
25 consisting of the VL, VH, C1 and CH1 domains; the Fd fragment consisting of the VH and CH1 domains; the Fv fragment consisting of the VL and VH domains of a single arm of an antibody; the dAb fragment which consists of a VH domain; isolated CDR regions and F(ab')₂ fragments, a
30 bivalent fragment including two Fab fragments linked by a disulphide bridge at the hinge region. Single chain Fv fragments are also included.

Humanised antibodies in which CDRs from a non-human
35 source are grafted onto human framework regions,

typically with the alteration of some of the framework amino acid residues, to provide antibodies which are less immunogenic than the parent non-human antibodies, are also included within the present invention and can be produced by the skilled person using techniques well known in the art.

A hybridoma producing a monoclonal antibody according to the present invention may be subject to genetic mutation or other changes. It will further be understood by those skilled in the art that a monoclonal antibody can be subjected to the techniques of recombinant DNA technology to produce other antibodies or chimeric molecules which retain the specificity of the original antibody. Such techniques may involve introducing DNA encoding the immunoglobulin variable region, or the complementarity determining regions (CDRs), of an antibody to the constant regions, or constant regions plus framework regions, of a different immunoglobulin. See, for instance, EP-A-184187, GB-A-2188638 or EP-A-0239400. Cloning and expression of chimeric antibodies are described in EP-A-0120694 and EP-A-0125023.

Hybridomas capable of producing antibody with desired binding characteristics are within the scope of the present invention, as are host cells, eukaryotic or prokaryotic, containing nucleic acid encoding antibodies (including antibody fragments) and capable of their expression. The invention also provides methods of production of the antibodies including growing a cell capable of producing the antibody under conditions in which the antibody is produced, and preferably secreted.

The antibodies described above may also be employed in the diagnostic aspects of the invention by tagging them

with a label or reporter molecule which can directly or indirectly generate detectable, and preferably measurable, signals. The linkage of reporter molecules may be directly or indirectly, covalently, e.g. via a peptide bond or non-covalently. Linkage via a peptide bond may be as a result of recombinant expression of a gene fusion encoding antibody and reporter molecule.

One favoured mode is by covalent linkage of each antibody with an individual fluorochrome, phosphor or laser dye with spectrally isolated absorption or emission characteristics. Suitable fluorochromes include fluorescein, rhodamine, phycoerythrin and Texas Red. Suitable chromogenic dyes include diaminobenzidine.

Other reporters include macromolecular colloidal particles or particulate material such as latex beads that are coloured, magnetic or paramagnetic, and biologically or chemically active agents that can directly or indirectly cause detectable signals to be visually observed, electronically detected or otherwise recorded. These molecules may be enzymes which catalyse reactions that develop or change colours or cause changes in electrical properties, for example. They may be molecularly excitable, such that electronic transitions between energy states result in characteristic spectral absorptions or emissions. They may include chemical entities used in conjunction with biosensors. Biotin/avidin or biotin/streptavidin and alkaline phosphatase detection systems may be employed.

In a further embodiment, the IPG antagonists are competitive antagonists, such as synthetic compounds or IPGs which are biologically inactive in the relevant respect, e.g. IPGs which do not have a TAF activity such

as enhancing the proliferation of tumour cells. An example of this latter type of antagonist is the use of porcine liver A-type IPGs in humans as described in GB-A-9828560.4 (WO00/38698) and related applications.

5 Synthetic compounds may be produced by chemical techniques or using combinatorial chemistry, and then screened for IPG antagonist activity. These compounds may be useful in themselves or may be used in the design of mimetics, providing candidate lead compounds for development as
10 pharmaceuticals. Synthetic compounds might be desirable where they are comparatively easy to synthesize or where they have properties that make them suitable for administration as pharmaceuticals, e.g. antagonist which are peptides may be unsuitable active agents for oral
15 compositions if they are degraded by proteases in the alimentary canal. Mimetic design, synthesis and testing is generally used to avoid randomly screening large number of molecules for a target property.

20 Pharmaceutical Compositions

IPGs or IPG antagonists can be formulated in pharmaceutical compositions. These compositions may comprise, in addition to one or more of IPGs or antagonists, a pharmaceutically acceptable excipient,
25 carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material may depend on the route of
30 administration, e.g. oral, intravenous, cutaneous or subcutaneous, nasal, intramuscular, intraperitoneal routes.

Pharmaceutical compositions for oral administration may be
35 in tablet, capsule, powder or liquid form. A tablet

may include a solid carrier such as gelatin or an adjuvant. Liquid pharmaceutical compositions generally include a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil.

5 Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included.

10 For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction, the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare
15 suitable solutions using, for example, isotonic vehicles such as sodium chloride injection, Ringer's injection, lactated Ringer's injection. Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.

20 Whether it is a polypeptide, antibody, peptide, small molecule or other pharmaceutically useful compound according to the present invention that is to be given to an individual, administration is preferably in a
25 "prophylactically effective amount" or a "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show benefit to the individual. The actual amount administered, and rate and time-course of
30 administration, will depend on the nature and severity of what is being treated. Prescription of treatment, e.g. decisions on dosage etc, is within the responsibility of general practitioners and other medical doctors, and typically takes account of the disorder to be treated,
35 the condition of the individual patient, the site of

delivery, the method of administration and other factors known to practitioners. Examples of the techniques and protocols mentioned above can be found in Remington's Pharmaceutical Sciences, 16th edition, Oslo, A. (ed), 1980.

The compositions may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated. Thus, in the treatment of cancer, the IPG antagonists can be administered in combination with other chemotherapy or radiotherapy.

Diagnostic Methods

Methods for determining the concentration of analytes in samples from individuals are well known in the art and readily adapted by the skilled person in the context of the present invention to determine whether an individual has an elevated level of IPGs, and so has or is at risk of cancer. The purpose of such analysis may be used for diagnosis or prognosis to assist a physician in determining the severity or likely course of the cancer and/or to optimise treatment of it. Examples of diagnostic methods are described in the experimental section below.

Preferred diagnostic methods rely on the detection of IPGs, employing biological samples such as blood, serum, urine or tissue samples (especially from a suspected tumour site).

The assay methods for determining the concentration of IPGs typically employ a binding agent having binding sites capable of specifically binding to the IPGs in preference to other molecules. Examples of binding

agents include antibodies (examples of which are provided above), receptors and other molecules capable of specifically binding IPGs. Conveniently, the binding agent is immobilised on solid support, e.g. at a defined location, to make it easy to manipulate during the assay.

The sample is generally contacted with a binding agent under appropriate conditions so that IPGs present in the sample can bind to the binding agent.

The fractional occupancy of the binding sites of the binding agent can then be determined either by directly or indirectly labelling the analyte or by using a developing agent or agents to arrive at an indication of the presence or amount of the analyte in the sample. Typically, the developing agents are directly or indirectly labelled (e.g. with radioactive, fluorescent or enzyme labels, such as horseradish peroxidase) so that they can be detected using techniques well known in the art. Directly labelled developing agents have a label associated with or coupled to the agent. Indirectly labelled developing agents may be capable of binding to a labelled species (e.g. a labelled antibody capable of binding to the developing agent) or may act on a further species to produce a detectable result. Thus, radioactive labels can be detected using a scintillation counter or other radiation counting device, fluorescent labels using a laser and confocal microscope, and enzyme labels by the action of an enzyme label on a substrate, typically to produce a colour change. In further embodiments, the developing agent or analyte is tagged to allow its detection, e.g. linked to a nucleotide sequence which can be amplified in a PCR reaction to detect the analyte. Other labels are known to those skilled in the art are discussed below. The developing agent(s) can be

used in a competitive method in which the developing agent competes with the analyte for occupied binding sites of the binding agent, or non-competitive method, in which the labelled developing agent binds analyte bound by the binding agent or to occupied binding sites. Both methods provide an indication of the number of the binding sites occupied by the analyte, and hence the concentration of the analyte in the sample, e.g. by comparison with standards obtained using samples containing known concentrations of the analyte.

The sample is generally contacted with the binding agent under appropriate conditions which allow the analyte in the sample to bind to the binding agent.

There is also an increasing tendency in the diagnostic field towards miniaturisation of such assays, e.g. making use of binding agents (such as antibodies or nucleic acid sequences) immobilised in small, discrete locations (microspots) and/or as arrays on solid supports or on diagnostic chips. These approaches can be particularly valuable as they can provide great sensitivity (particularly through the use of fluorescent labelled reagents), require only very small amounts of biological sample from individuals being tested and allow a variety of separate assays can be carried out simultaneously. This latter advantage can be useful as it provides an assay employing a plurality of analytes to be carried out using a single sample. Examples of techniques enabling this miniaturised technology are provided in WO84/01031, WO88/1058, WO89/01157, WO93/8472, WO95/18376/ WO95/18377, WO95/24649 and EP 0 373 203 A. Thus, in a further aspect, the present invention provides a kit comprising a support or diagnostic chip having immobilised thereon a plurality of binding agents, at least one of which is

capable of binding to an IPG as described above,
optionally in combination with other reagents (such as
labelled developing reagents) needed to carrying out an
assay.

Materials and Methods

Anti-IPG Antibodies

Anti-IPG antibodies were produced by culturing hybridoma
cell lines 2F7, 2D1 and 5H6 deposited at European
Collection of Cell Cultures (ECACC) under accession
numbers 98051201, 98031212 and 98030901, and isolating
the anti-IPG antibodies thus produced.

Cellulose Purification of IPGs

Microcrystalline cellulose (Merck Avicel®) was fined by
suspension in double-distilled water followed by standing
for 10 minutes and decanting of the unsettled suspension.
After 12 repeats, the settled material was shaken
vigorously with two volumes of fresh water and poured
into the top of a Pharmacia® glass column of 16 mm i.d.
with the top plunger removed. Water was allowed to flow
out under gravity at 3 ml/min. After allowing the
solvent level to drop to the top of the packed bed, the
plunger was replaced and water pumped through the column
at 5 ml/min using a Pharmacia® peristaltic pump. The
column was treated with the following solvents:

Water	150ml
50% aqueous methanol	50ml
Methanol	50ml
Butanol/methanol/water (4:1:1)	150ml

All solvents were degassed prior to use by brief swirling
under vacuum. After the final solvent, the plunger was
removed, and the solvent level allowed to drop to the top

of the packed bed under gravity, whereupon flow was stopped.

The sample (2 ml of conditioned cell medium or IPG extract from two rat livers) was swirled for 5 minutes with 2 ml of the cellulose suspension and the entire mixture lyophilised.

The dry cellulose material was suspended in 2ml of butanol/ethanol/water (B:W:E) (4:1:1) and applied gently and evenly to the top of the packed bed. The remaining material was washed in using a further 2ml of the solvent. The solvent level was again dropped to the bed level before commencing pumping with B:W:E (4:1:1) at 5 ml/min. Fractions were collected in glass vessels as follows, either by hand into flasks or using a Gilson model 204 fraction collector set to 1 minute/tube:

Butanol/methanol/water (4:1:1)	150ml
Methanol	50ml
50% aqueous methanol	10 x 5ml fractions
Water	20 x 5ml fractions
HCl(aq.) pH 1.3	10 x 5ml fractions
HCl(aq.) pH 1.3	75ml

The samples were evaporated to dryness using a rotary evaporator or Speedvac at below 30°C. Samples were dissolved in 200µl of water and stored at -20°C.

Preparation of conditioned medium (CM) from H4IIE cells

(a) H4IIE cells were cultured on 75 cm² Falcon cap vented tissue culture flasks at cell density of $2-6 \times 10^4$ cells/cm² using DMEM (Gibco Cat. No. 31885), supplemented at 5% with a 1:1 mixture of foetal calf serum and heat inactivated calf serum and incubate at

37°C, 5% CO₂ atmosphere until 100% confluence.

(b) At confluence (15-17 x10⁶ cells/T75 flask), the cells were washed twice using 15-20 mL Hank's Balance Salt Solution pH 7.4 (HBSS, Gibco Cat No. 14174-053) and 5ml Trypsin-EDTA solution (Sigma Cat. No. T 4171) was added. The excess of Trypsin was then removed, leaving in the flask only 2 ml of liquid which was then incubated for 10 min.

(c) When the cells detached, 8 ml of growth medium was added and the mixture was centrifuged for 5 min at 1000 rpm. The supernatant was discharged and the pellet resuspended by adding 10 ml growth medium. The cell suspension was transferred into a sterile 25 ml universal container, 10 ml of the same medium was added, the cap closed tightly and the container rocked several times to produce a homogeneous cell suspension.

(d) The cells were subcultured 1/10 or 1/20 (1-2.5 x 10⁴/cm²) in 75 cm² Falcon cap vented flask, using the same medium and incubation conditions described in step (a). When the cells seeded at 1/10 reached 70% confluence (11.5-12.0 x 10⁶ cells/ T75 flask, normally 3 days after), the culture was washed three times with 20 ml serum free DMEM and then incubated for 3 hours. This procedure was repeated twice.

(e) The culture was incubated for 48 hr at 37°C, 5% CO₂ atmosphere, and then the conditioned medium was transferred into a centrifuge tube, and cell debris removed by centrifugation at 1000g for 10 min at 4°C and then at 105,000 g for 60 min at 4°C. The supernatant was sterilized by filtration and stored at -80°C.

Biosynthetic labelling and extraction of conditioned medium

(a) H4IIE cells grown to 50-60% confluence in T75 cap-vented flask Falcon (Falcon Cat. No F3111) ($8-9 \times 10^6$ cells/T75 flask) in DMEM medium, supplemented with 5% foetal calf serum and 5% heat inactivated calf serum, were washed twice using HBSS at 37°C. 10 mL of fresh growth medium was added and then 250 mCi of the ^3H -labelled precursors (Glucosamine, Galactose, Mannose and myo-Inositol). The culture was incubated at 37°C, 5% CO_2 atmosphere for 24 hours.

(b) After 24 hours, the cells were washed three times as described in step (d) above and the cells incubated and the medium harvested as in step (e).

(c) The supernatants obtained were adjusted to pH 3.0 with HCl (c), and prewashed activated charcoal (25 mg/mL) added and stirred at 4°C for 30 min. The suspension was transferred into centrifuge tubes and the charcoal spun down by centrifugation at 20,000 g for 30 min at 4°C.

(d) The supernatant was sterilized using a 0.2 μm pore size filter and stored at 4°C. If it was not required for immediate use, it was freeze-dried and kept at -80°C until needed.

Bioactivity assay of H4IIE conditioned medium by measuring the proliferation of FaO cells

(a) FaO cells grown at 60-70% confluence in T75 Falcon cap-vented tissue culture flasks Gibco Cat.No F3111 with RPMI-1640 (Gibco Cat. No. 21875), supplemented at 5% with a 1:1 mixture of foetal calf serum and heat inactivated calf serum were washed twice with 20 ml of HBSS pH 7.4

(Gibco Cat. No. 14174). The cells were detached with Trypsin-EDTA solution (Sigma Cat. No. T4171) as described in steps (b) and (c) of "Preparation of conditioned medium from H4IIE" protocol set out above.

(b) The cell suspension was counted twice and the required volume to dilute the suspension at a density of $2.5-3.0 \times 10^3$ cell/ml was calculated. The calculated volume of growth medium was added and the cells were transferred into a sterile 25 ml universal container. The container was rocked gently several times to produce a homogenous cell suspension.

(c) The cells were plated in a 96-wells Falcon U bottom shape microwell plate by adding 100 μ l of cell suspension per well and incubated at 37°C in a 5% CO₂ atmosphere incubator. After 24 hours, each well was washed three times with 100 μ l of serum free medium and incubates for a further 24 hours.

(d) The medium was replaced by adding serum free medium and the cells incubated for three additional hours. Meanwhile, the IPG samples or medium supplementation at different concentrations (ranging from 1% to 20.0%) of H4IIE conditioned medium were prepared. 100 μ l serum free medium was added to the controls, 100 μ l completed growth medium was added to the positive controls and 100 μ l of medium supplemented with H4IIE conditioned medium.

(e) The cells were incubated at 37°C in a 5% CO₂ atmosphere incubator for 18 hours and then 1 mCi ³H-thymidine was added to each well. After 4 hours incubation, the radioactive medium was removed and replaced with 50 μ l trypsin-EDTA solution. The cells were incubated for 10 min and the DNA collected on

filters (Helis Bio Ltd. Cat No 11731) using a cell harvester (SKATRON 12 well/cell harvester). The filters were transferred to scintillation vials, 2 ml of scintillation cocktail was added to each vial and counted for one minute in a Beckman scintillation counter.

Assays

Phosphate analysis and PDH activation assays were carried out as described in Caro et al, 1997.

Results

Fractionation of conditioned medium by cellulose column chromatography

Conditioned medium biosynthetically labelled with ^3H myo-inositol (squares), glucosamine (circles) or galactose (open circles) at $1.35\text{E}5$, $2.85\text{E}5$ and $2.50\text{E}5$ cpm respectively, obtained as described in Methods, was separated by cellulose column chromatography, 5 ml fractions were collected and 200 μl counted in a Beckman liquid scintillation counter. The results are presented in Figure 1 as counts per minute (cpm) per fraction. The arrows indicate the solvent system used at a particular fraction. The results in Figure 1 show that these radioactive components are incorporated into TAF fractions obtained from the supernatant and that the TAF fractions can be isolated by the same procedures as IPGs.

Effects on DNA synthesis of conditioned medium

Conditioned medium at two different concentrations (10% and 20%) was assayed for stimulating activity in test Fao cells, as under Methods. The fractions obtained by purification of conditioned medium (2 ml) by cellulose column chromatography were concentrated to dryness, dissolved in water (200 μl), and also assessed for

activity (4 μ l). The results are presented in Figure 2 as radioactivity (cpm) obtained after incorporation of 3 H-thymidine into DNA, for control (no serum addition), FCS (5% foetal calf serum + 5% foetal bovine serum addition), and the fractions obtained. The arrows indicate the eluent system used at a particular fraction. The results show that some of the fractions, and in particular fraction 21 causes proliferation of Fao cells as does the addition of conditioned medium.

Phosphate analysis of conditioned medium

Conditioned medium (2 ml) was fractionated by column chromatography using cellulose as described under methods. Fractions (5 ml) were collected, concentrated to dryness, dissolved in water (200 μ l) and a portion (10 μ l) was assayed for phosphate as described in Caro et al, 1997. Briefly, samples were evaporated to dryness and hydrolized with perchloric acid (70% by volume) at 180°C for 30 min. After cooling at room temperature, distilled water (250 μ l) was added. Ammonium molybdate and ascorbic acid were sequentially added, yielding final concentrations of 3 and 72 mM, respectively. Colour development was achieved by heating at 95°C for 15 min. Optical absorbance was measured at 650 nm. The results are presented in Figure 3 as OD obtained per fraction. Arrows indicate the eluent system used at a particular fraction. The results show that the fractions co-purifying with TAF activity contain phosphate, supporting the fact that TAFs are IPGs.

Effect on the proliferation of FaO cells of conditioned medium treated with anti-IPG monoclonal antibodies

Test FaO cells were used to measure the stimulating activity of conditioned medium as described in methods, in the presence of the anti-IPG monoclonal antibodies

2D1, 5H6 and 2F1 at a concentration of 1 µg per well. The results in Figure 4 are presented as radioactivity (cpm) obtained after incorporation of ³H-thymidine into DNA, for control (no serum addition), FCS (5% foetal calf serum + 5% foetal bovine serum addition) and samples containing the antibodies. The results show that all three antibodies antagonise the proliferating activity caused by the addition of conditioned medium to the Fao cells, and that antibody 2F1 had the greatest inhibitory effect.

Rat liver IPGs A and P fractionation profiles

The IPGs obtained from two rat livers were fractionated using cellulose column chromatography as described in Methods. The fractions obtained were evaporated to dryness, redissolved in water (200 µl), and a portion of the solution (1.25 µl) assayed for biological activity in triplicates. The results are shown as radioactivity (cpm) obtained after incorporation of ³H-thymidine into DNA of 3T3EFGTR17 cells, for control (no serum addition), FCS (10% foetal calf addition), and fractions. All fractions were assayed at a final concentration of 1/80, except for fraction 26 of IPG-P (Figure 5) which was also tested at a final concentration of 1/40 and 1/160. All fractions of the IPG-A (Figure 6) were tested at a final concentration of 1/80. The arrows indicate the eluent system used at a particular fraction. The results indicate that cellulose chromatography is an effective technique for the isolation and purification of IPGs, e.g. IPGs obtained from natural source materials.

PDH activity of Fraction 26 of IPG-P type

PDH activity was measured as described in Caro et al, 1997. A portion (10 ml) of fractions 22, 26 and 27 was assayed. The result is presented as activation of the PDH complex expressed as units per gram of tissue. One

unit of IPG PDH activity is the amount required to increase the basal rate of NADH production by 50%.

Fraction 22 = 0.123 units/g

Fraction 26 = 2.09 units/g

Fraction 27 = 0.523 units/g.

The results show that the IPGs purified from rat liver using cellulose chromatography have a characteristic P-type IPG biological activity.

Phosphate analysis of rat liver IPG-P

The IPG-P obtained from two rat livers was fractionated by column chromatography using cellulose as described under Methods. Fractions (5 ml) were collected, concentrated to dryness, dissolved in water (200 µl) and a portion (100 µl) was assayed for phosphate as described in Caro et al, 1997. Each fraction was evaporated to dryness and hydrolyzed with perchloric acid (70% by volume) at 180°C for 30 min. After cooling at room temperature, distilled water (250 µl) was added. Ammonium molybdate and ascorbic acid were sequentially added, yielding final concentrations of 3 and 72 mM, respectively. Colour development was achieved by heating at 95°C for 15 min. Optical absorbance was measured at 650 nm. The results in figure 8 are presented as OD obtained per fraction. Arrows indicate the eluent system used at a particular fraction. The results show that the P-type IPGs purified using cellulose chromatography contain phosphate.

Deposits

The deposit of hybridomas 2F7, 2D1 and 5H6 in support of this application was made at the European Collection of Cell Cultures (ECACC) under the Budapest Treaty by

Rademacher Group Limited (RGL), The Windeyer Building, 46
Cleveland Street, London W1P 6DB, UK. The deposits have
been accorded accession numbers accession numbers
98051201, 98031212 and 98030901 on 12 May 1998 ('201) and
5 9 March 1998 ('212 and '901). RGL give their unreserved
and irrevocable consent to the the materials being made
available to the public in accordance with appropriate
national laws governing the deposit of these materials,
such as Rules 28 and 28a EPC. The expert solution under
10 Rule 28(4) EPC is also hereby requested.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
219

References:

The references referred to herein are expressly incorporated by reference.

5 WO98/11116 and WO98/11117 (Rademacher Group Limited).

Rademacher et al, Brazilian J. Med. Biol. Res., 27:327-341, 1994.

10 Caro et al, Biochem. Mol. Med., 61:214-228, 1997.

Witters et al, J. Bio. Chem., 263:8027-8036, 1986.

15

methanol.

Claims:

1. Use of a substance which is an inositolphosphoglycan (IPG) antagonist having the property of reducing tumour cell proliferation for the preparation of a medicament for the treatment of cancer.

2. The use of claim 1, wherein the IPG antagonist is:
(a) a substance which is capable of inhibiting the release of IPGs; or,

(b) a substance capable of reducing the levels of IPGs by binding to the IPGs; or,

(c) a substance which is a competitive agent which capable of reducing an effect of IPGs.

3. The use of claim 2, wherein the antagonist is a competitive IPG antagonist.

4. The use of claim 2, wherein the IPG antagonist is an anti-IPG antibody which is capable of specifically binding IPGs.

5. The use of claim 4, wherein the antibody capable of neutralising an activity of the IPGs.

6. The use of claim 5, wherein activity of the IPGs is the proliferation of tumour cells.

7. The use of any one of claims 4 to 6, wherein the antibody is a monoclonal antibody produced by hybridoma 2F7, 2D1 or 5H6, deposited at ECACC under accession numbers 98051201, 98031212 and 98030901.

8. The use of claim 2, wherein the antagonist is an inhibitor of glycosylphosphatidylinositol specific phospholipase type C (GPI-PLC).

9. Use of the presence or amount of inositolphosphoglycans (IPGs) in a sample from a patient for the diagnosis and/or prognosis of cancer.

10. A method for the diagnosis and/or prognosis of cancer, the method comprising determining the presence or amount of inositolphosphoglycans in a sample from a patient.

11. The method of claim 10, wherein the presence or amount of the IPGs is causing tumour cell proliferation.

12. The method of claim 10 or claim 11, wherein the method comprises the steps of:

(a) contacting a sample from a patient with a solid support having immobilised thereon a binding agent having binding sites which are capable of specifically binding to the IPGs with a sample from a patient under conditions in which the IPGs bind to the binding agent; and,

(b) determining the presence or amount of the IPGs bound to the binding agent.

13. The method of claim 12, wherein step (b) comprises (i) contacting the solid support with a developing agent which is capable of binding to occupied binding sites, unoccupied binding sites or the bound IPGs, the developing agent comprising a label and (ii) detecting the label to obtain a value representative of the presence or amount of the IPGs in the sample.

14. The method of claim 13, further comprising comparing the value with standards from healthy or cancerous tissues.

15. The method of claim 13 or claims 14, wherein the

label is a radioactive label, a chemiluminescent label, a fluorophor, a phosphor, a laser dye, a chromogenic dye, a macromolecular colloidal particle, a latex bead which is coloured, magnetic or paramagnetic, or an enzyme which catalyses a reaction producing a detectable result.

15. The method of any one of claims 12 to 15, wherein the binding agent immobilised on the solid support is an antibody which is capable of binding to the IPGs.

16. The method of any one of claims 12 to 16, wherein the binding agent is immobilised at a predefined location on the solid support.

17. Use of microcrystalline cellulose for purifying or isolating a P or A-type substance, wherein the substance is a cyclitol containing carbohydrate which is:

- (i) a P-type substance having the biological activity of activating pyruvate dehydrogenase (PDH) phosphatase; or,
- (ii) an A-type substance having the biological activity of inhibiting cAMP dependent protein kinase.

18. The use of claim 17, wherein the use involves contacting a sample containing P or A-type substance with a column containing cellulose and eluting the substance from the column.

19. A method of purifying or isolating a P or A-type substance, wherein the substance is a cyclitol containing carbohydrate which is:

- (i) a P-type substance having the biological activity of activating pyruvate dehydrogenase (PDH) phosphatase; or,
- (ii) an A-type substance having the biological


activity of inhibiting cAMP dependent protein kinase;
wherein the method comprises:

- (a) loading a column containing microcrystalline cellulose with a sample containing the P or A-type substance so that P or A-type substance binds to the column; and,
- (b) eluting the P or A-type substance from the column.


20. The method of claim 19, further comprising the step of dissolving the sample containing the P or A-type substance in 4/1/1 butanol/water/ethanol (B:W:E) prior loading on the column.

21. The method of claim 19 or claim 20, further comprising the step of washing the column with B:W:E and methanol.

1/10




2/10



3/10


FIGURE 2
Proliferation assay - FaO cells



4/10


FIGURE 3

H4 Conditioned Medium Fractionation - Phosphate Analysis




5/10

FIGURE 4
Effects of mAb on H4IIE Conditioned Medium
Cell proliferation (FaO cells)




6/10

FIGURE 5
Proliferation Assay - Liver IPG P - Fibroblasts cells




7/10

FIGURE 6 A
Proliferation Assay - Liver IPG A - Fibroblasts cells




8/10

FIGURE 6B
Proliferation Assay - Liver IPG A - Fibroblasts cells




9/10

FIGURE 7A
Liver IPG P - Cellulose Purification- Phosphate Analysis



10/10

FIGURE 7B
Liver IPG P - Cellulose Purification- Phosphate Analysis



Attorney Docket No.: 1012E-000300US
 Client Reference No.: SJK/FP5938444

DECLARATION

As a below named inventor, I declare that:

My residence, post office address and citizenship are as stated below next to my name; I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: **TREATMENT AND DIAGNOSIS OF CANCER USING INOSITOLPHOSPHOGLYCANS ANTAGONISTS** the specification of which is attached hereto or was filed on June 22, 2001 as Attorney Docket No. 1012E-000300US, and was amended on (if applicable).

I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56. I claim foreign priority benefits under Title 35, United States Code, Section 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Country	Application No.	Date of Filing	Priority Claimed Under 35 USC 119
PCT	PCT/GB99/04382	December 23, 1999	Yes
GB	9828564.6	December 23, 1998	Yes

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below:

Application No.	Filing Date



I claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application No.	Date of Filing	Status
PCT/GB99/04382	December 23, 1999	Inactive

Attorney Docket No.: 1012E-000360US
 Client Reference No.: S/K/EP5938444

Full Name of Inventor 1:	Full Name: RADEMACHER, Thomas William		
Residence & Citizenship:	City: Oxford	State/Region/Country: United Kingdom	Country of Citizenship: United States
Post Office Address:	Post Office Address: Foxcombe, The Ridgeway Boats Hill, Oxford OX1 5QU United Kingdom		
Full Name of Inventor 2:	Full Name: CARO, Hugo		
Residence & Citizenship:	City: LONDON	State/Region/Country: United Kingdom	Country of Citizenship: Argentina
Post Office Address:	Post Office Address: 52 TURNELL PARK RD, LONDON N7 0DT		
Address:	Address: 44 Bedford Avenue, Bedford, Hertfordshire AL5 2EP United Kingdom		

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 1:	Signature of Inventor 2:
	
RADEMACHER, Thomas William	CARO, Hugo
Date 20/06/01	Date 21/06/01